

Ultimate barricade against pre-emergent weeds that is exempt from poison scheduling




syngenta®

BARRICADE Turf Herbicide controls a wide range of pre-emergent grassy weeds and importantly breaks the germination cycle of *Poa annua*. You can trust BARRICADE to last for up to 6 months so your application window is more flexible.

BARRICADE binds high in the soil profile ensuring turf roots below the barrier are not affected and your turf stays in prime condition.

What BARRICADE offers you:

- Up to six months control with a single application per season with little risk of herbicide to your turf
- An economical solution for many grassy weeds including African Lovegrass, Parramatta Grass, Summer Grass, Crab Grass, Crowsfoot Grass and Winter Grass
- Flexible application window – well before weeds germinate
- Exempt from poison scheduling
- Low odour, non-staining formulation with the flexibility to delay wash-in for a few days if necessary
- A complete couch solution for *Poa annua* control when mixed with MONUMENT LIQUID Turf Herbicide



BARRICADE can be used on the following turf species:

Warm Season Turf Grasses

Bahia Grass, Buffalo Grass, Carpet Grass, Common Couch, Hybrid Couch, Kikuyu, Queensland Blue Couch, Seashore Paspalum, Zoysia

Refer to the registered label for full details.

SpraySURE | Accuracy and Performance

To achieve great results, accurate application is important.

Target	Syngenta Nozzle	Optimal Water Rate	Optimal Pressure Range
Root and soil So	XC 08	> 500 L/ha	2 to 4 bar

BARRICADE can deliver up to 6 months residual control. Positioning of applications can be carried out well in advance of weeds germinating and still deliver season long control.

Example: Spring applications for the control of summer grassy weeds like *Digitaria* (Summer Grass and Crab Grass) and *Eleusine* (Crowsfoot Grass) should ideally be applied around the end of August. Control will last up to the end of February. Autumn applications for the control of *Poa* (Winter Grass) should ideally be applied around the end of February. Control will last up to the end of August.

BARRICADE as a pre-emergent herbicide works in the top soil layers. Application targeted towards delivering BARRICADE close to the soil surface is best suited to ensure optimum results. BARRICADE needs to be washed in within 7 days after application.

Target Weeds	Rate	SpraySure
Barnyard Grass, Crab Grass, Crowsfoot Grass, Parramatta Grass, Rat's Tail Fescue, Summer Grass	1 to 3 L/ha	So
African Lovegrass, Bahia Grass, Kentucky Blue Grass	2 to 3 L/ha	So
Paspalum	2 to 4 L/ha	So
Winter Grass	1 to 4 L/ha	So

Refer to the registered label for full details and instructions for use. **F** = Foliar applied **So** = Soil applied **Su** = Surfactant required

Active Ingredient

BARRICADE contains 480 g/L proflamifen, a member of the dinitroaniline herbicides

GROUP **D** HERBICIDE

Formulation and Packaging

5 L pack size

A suspension concentrate (SC) with minimal odour and non-staining properties. BARRICADE mixes completely with water and produces a homogenous easy to apply suspension.

BARRICADE can be tank mixed with MONUMENT LIQUID. Observe the mixing instruction for MONUMENT LIQUID with regards to water pH and the addition of a non-ionic surfactant (buffer solution prior to MONUMENT LIQUID addition to ~pH 7 and add a non-ionic surfactant at 0.25% v/v).



www.greencast.com.au

 **Barricade**[®]
Herbicide

syngenta[®]

For more information and to download the full registered label and MSDS, ask your Syngenta Agent or go to www.greencast.com.au

Syngenta Australia Pty Limited, Level 1, 2-4 Lyonpark Road, Macquarie Park NSW 2113. ABN 33 002 933 717.

® Registered trademark of Syngenta Group Company. ™ Trademark of a Syngenta Group Company.

All products written in uppercase are registered trademarks of a Syngenta Group Company. ©Syngenta 2015 SB 15/109/R