



POSTERITY® Fungicide holds strong for many moons.



syngenta®

NEW registration for Spring Dead Spot and Microdochium Patch.


Ideally suited for applications to sporting fields and golf course fairways, POSTERITY® Fungicide is the latest generation SDHI in the turf market. Providing exceptional Spring Dead Spot, Dollar Spot, and Microdochium Patch control, it sets a new standard holding stronger for many moons.

What POSTERITY® offers:

- A step-change for fairway and sportsfield Spring Dead Spot control compared to traditional methods, providing unprecedented control at extremely low rates.
- Powerful, preventative Spring Dead Spot control that stays where you place it in the soil profile, working longer and harder.
- Curative and preventative control against Dollar Spot and Microdochium Patch.
- A formulation that is compatible with a wide range of commonly used products including PRIMO® MAXX so you can supercharge your tank mix.
- Balanced distribution within the plant and outstanding rainfastness – it binds quickly and strongly to the plant's wax layer and then slowly distributes into the plant tissue.
- Smarter chemistry that is exempt from poison scheduling.
- A highly effective option at rates as low as 500 mL/ha to 1 L/ha.

POSTERITY®: A step change in Spring Dead Spot Control

Spring Dead Spot development post spring green-up following two autumn applications of POSTERITY®



Dedicate® is a Registered Trademark of the Bayer Group. Trial Reference: WR-FPSF055C-2020AU-SYN482.

To achieve great results, accurate application is important

Target	Syngenta Nozzle	Optimal Water Rate	Optimal Pressure Range
Leaf and Crown F	XC25 or XC04	350-800 L/ha	2-4 bar
Root So	XC08	1000 L/ha	2-4 bar

For preventative Spring Dead Spot control, apply prior to symptom development in mid-Summer to Autumn and use the higher rate when historical disease pressure was high. Ensure application is irrigated with approximately 6 mm of water directly after application.

To combat Dollar Spot, start the season strong with an early application of POSTERITY® when conditions are favourable for disease outbreak or as an early curative application at first signs of disease. Favourable conditions include: temperatures between 15°C and 30°C, continuous high humidity, cool nights and intense dews.

For Winter Fusarium/Microdochium Patch control, use POSTERITY® as a rotation partner.



UNTREATED DOLLAR SPOT



POSTERITY® 0.5 L/ha (2 applications)

Trial ref: Clark2016_RU

Target Diseases	Rate/ha	SpraySure
Dollar Spot, Winter Fusarium/ Microdochium Patch	0.5 to 1 L	F
Spring Dead Spot	0.5 to 1 L	So

Refer to the registered label for full details and instructions for use.

F = Foliar applied **So** = Soil applied

Fungicide Group

POSTERITY® is powered by ADEPIDYN™ technology; a member of the carboxamide (SDHI) fungicide group.

GROUP **7** FUNGICIDE

Formula & Packaging

500 mL pack size.

Suspension concentrate (SC) that mixes completely with water and may be tank mixed with many commonly used fungicides, insecticide, herbicides and fertilisers.



Posterity®
Turf fungicide

syngenta®

For more information, ask your Syngenta Agent or call Syngenta Customer Service on 1800 022 035 or visit syngentaturf.com.au