

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Scarlet Trio

ADVANCED INSECTICIDE

**ACTIVE CONSTITUENTS: 116 g/L CLOTHIANIDIN
83 g/L CLOFENTEZINE
13 g/L ABAMECTIN**

GROUP 4A 10A 6 INSECTICIDE

**For the control of billbug, African black beetle, Argentinian scarab,
Argentine stem weevil, couch fly, couch mite, ground pearl, mole cricket
in turf as specified in the Directions for Use table**

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING



Indigo Specialty Products Pty Ltd
ABN 15 631 459 660
3/49 Donaldson Road,
Rocklea QLD 4106
Ph: 0402 735 887

**Contents
5 L**



Scarlet Trio

ADVANCED INSECTICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. The container can be recycled if it is clean, dry, free of visible residues and has the *drumMUSTER* logo visible. Triple-rinse container for disposal. Dispose of rinsate or any undiluted chemical according to state legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any *drumMUSTER* collection or similar container management program site. The cap should not be replaced, but may be taken separately. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. May irritate the eyes and nose and throat. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. Do not inhale spray mist. When opening the container and preparing the product for use wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves and a disposable mist face mask covering mouth and nose. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing). Wash hands after use. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126, New Zealand 0800 764 766.

SDS

Additional information is listed in the safety data sheet (SDS). A safety data sheet for ProForce SCARLET TRIO Advanced Insecticide is available from Indigo Specialty Products Pty Ltd on request.

May cause damage to the central nervous system and the peripheral nervous system through prolonged or repeated exposure.

Do not eat, drink or smoke when using this product.
Get medical advice/attention if you feel unwell.

CONDITIONS OF SALE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Indigo Specialty Products Pty Ltd or under abnormal conditions. Indigo Specialty Products Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

IN A TRANSPORT
EMERGENCY

DIAL 000

POLICE OR FIRE BRIGADE

SPECIALIST ADVICE IN AN EMERGENCY ONLY

0417 653 654

ALL HOURS – AUSTRALIA WIDE



5L

Batch No.

Date of
Manufacture

APVMA Approval No.: 92166/138402



ADVANCED INSECTICIDE

DIRECTIONS FOR USE

Restraints

- DO NOT use on fine textured compacted clay soils without amelioration prior to treatment application.
- DO NOT apply if heavy rains or storms are forecast within 48 hours.
- DO NOT irrigate to the point of field runoff for at least 48 hours after application.
- DO NOT apply if rainfall imminent. The effect of this product could be diminished if rain falls within 6 hours of application.
- DO NOT apply to turf under heat or moisture stress.
- DO NOT apply by backpack.
- DO NOT apply by aircraft.
- DO NOT apply by vertical sprayer.
- DO NOT apply through any type of irrigation system or ultra low volume spray system.
- DO NOT apply more than 2 sprays per season in turf.
- DO NOT apply with a nozzle height greater than 50 cm above the ground.

SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift. DO NOT allow bystanders to come into contact with the spray cloud. DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone table/s below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas. DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application. DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

DO NOT apply by a boom sprayer unless the following requirements are met:

- Spray droplets are not smaller than a COARSE spray droplet size category.
- Minimum distances between the application site and downwind sensitive areas are observed (see the following table titled 'Buffer zones for boom sprayers').

Buffer zones for boom sprayers

Application rate	Boom height above the target canopy	Mandatory downwind buffer zones			
		Bystander areas	Natural aquatic areas	Pollinator areas	Vegetation areas
3 L/ha	0.5 m or lower	0 metres	10 metres	110 metres	0 metres

Situation	Pest	Rate	Critical Comments
Turf (including Golf Courses, Sportsfields, and other sport and recreational turf areas)	Billbug (La Plata Weevil) (<i>Sphenophorus brunnipennis</i>)	3 L/ha	Monitor adult activity through late spring and early summer. Spray when numbers peak or when small larvae (4 mm) are found in the thatch or surface soil typically around late November to early December. Early application is essential to minimise damage to turf due to feeding. Ensure product placement as close to soil surface as possible. Preferably spray onto wet or dewy turf. Irrigate with at least 3 mm of water commencing within 1 hour of application. DO NOT irrigate to the point of runoff. A second application may be required to provide control.
	Argentine Stem Weevil (<i>Listronotus bonariensis</i>), Argentinian Scarab (<i>Cyclocephala signaticollis</i>), Mole Cricket (<i>Gryllotalpa</i> sp.)		Monitor adult activity through spring and early summer. Spray when peak numbers (preventatively) or when first visual symptoms are observed (curatively) typically around late September to January. Early application is essential to minimise damage to turf due to feeding. Argentinian Scarab: A second application may be required to provide control.
	African Black Beetle (<i>Heteronychus arator</i>)		Apply at peak egg hatch (African Black Beetle – late September through to mid-November), or when small larvae are present. Ensure product placement as close to soil surface as possible. Preferably spray onto wet or dewy turf. Irrigate with at least 3 mm water commencing within 1 hour of application. DO NOT irrigate to the point of runoff. A second application may be required to provide control.
	Ground pearl (<i>Margoroides</i> spp.)		Apply when adults or crawlers are detected generally from early spring to the end of summer. Apply in 400 – 800 L/ha.
	Couch Mite (<i>Aceria cynodoniensis</i>), Couch Fly (<i>Delia urbana</i>)		Apply product in an early curative situation (after first symptoms are apparent). Apply in 400-800 L of water per hectare. Best results are achieved if applied as populations begin to build rather than at the peak of population growth. A second application may be required to provide control.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

TURF: DO NOT GRAZE TREATED TURF/LAWN OR FEED TURF/LAWN CLIPPINGS FROM ANY TREATED AREA TO POULTRY OR LIVESTOCK

GENERAL INSTRUCTIONS

ProForce SCARLET TRIO ADVANCED Insecticide is a combination insecticide providing contact and residual control of a wide range of insect pests in turf. Clothianidin, clofentezine and abamectin have different modes of action so together provide long term activity on a wide spectrum of pests.

MIXING

Use only clean spraying equipment in good condition. Half fill the spray tank with clean water and commence agitation. Add the required quantity of product directly to the water,

then add mixing partner and complete filling to the final volume. Spray immediately and maintain agitation throughout. DO NOT allow spray mixture to remain in the tank overnight or for long periods without agitation.

APPLICATION

Apply by ground boom sprayer, low pressure hand wand or hand gun sprayer. To be effective SCARLET TRIO ADVANCED requires thorough spray coverage. Ensure that equipment is properly calibrated to give an even distribution at the correct volume. Application volume should be adequate to ensure thorough and even coverage of turf

leaves with penetration into the crowns. Total application volume should be 400 to 800 L/ha. Use coarse droplets (e.g. Air Induction flat fan 025 to 04 nozzles). In higher cut turf (>15 mm) a significant spray shielding effect can occur, impacting negatively on spray penetration and even coverage at low application volumes.

INSECTICIDE RESISTANCE WARNING

For insecticide resistance management ProForce SCARLET **GROUP 4A 10A 6 INSECTICIDE** TRIO ADVANCED Insecticide is a combination of Group 4A, Group 10A and Group 6 insecticide. Some naturally occurring insect biotypes resistant to SCARLET TRIO ADVANCED and other Group 4A, Group 10A or Group 6 insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if SCARLET TRIO ADVANCED or other Group 4A, Group 10A or Group 6 insecticides are used repeatedly. The effectiveness of SCARLET TRIO ADVANCED on resistant individuals could be significantly reduced. Since the occurrence of resistant insects is difficult to detect prior to use, Indigo Specialty Products Pty Ltd accepts no liability for any losses that may result from the failure of SCARLET TRIO ADVANCED to control resistant insects. SCARLET TRIO ADVANCED may be subject to specific resistance management strategies. For further information contact your local supplier, Indigo Specialty Products representative or local agricultural department agronomist.

RE-ENTRY PERIOD

General Public

Do not allow entry into treated areas until the spray has dried.

Occupational Users

Do not allow entry into treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF HONEY BEES AND OTHER INSECT POLLINATORS

Highly toxic to bees. Harmful to bee brood. Clothianidin has systemic action. DO NOT spray while bees are actively foraging. DO NOT allow spray drift to flowering weeds or flowering crops in the vicinity of the treatment area. Before spraying, notify beekeepers to move hives to a safe location with an untreated source of nectar and pollen, if there is potential for managed hives to be affected by the spray or spray drift. Risk to bees is reduced by spraying early morning and late evening while bees are not foraging. Residues may remain at levels toxic to bees for several days following application.

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. May irritate the eyes and nose and throat. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. Do not inhale spray mist. When opening the container and preparing the product for use wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves and a disposable mist face mask covering mouth and nose. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing). Wash hands after use. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126, New Zealand 0800 764 766.

SDS

Additional information is listed in the safety data sheet (SDS). A safety data sheet for ProForce SCARLET TRIO ADVANCED Insecticide is available from Indigo Specialty Products Pty Ltd on request. Call Customer Service on 02 9389 2455.

CONDITIONS OF SALE:

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Indigo Specialty Products Pty Ltd or under abnormal conditions. Indigo Specialty Products Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

Integrated Pest Management

Toxic to beneficial arthropods. Minimise spray drift to reduce harmful effects on beneficial arthropods outside the treatment area.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DRIFT WARNING - DO NOT apply under weather conditions or from spraying equipment which could be expected to cause spray to drift onto adjacent, crops, crop lands, pastures or livestock.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight.

1 L

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

DrumMUSTER

The container can be recycled if it is clean, dry, free of visible residues and has the *drumMUSTER* logo visible. Triple-rinse container for disposal. Dispose of rinsate or any undiluted chemical according to state legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any *drumMUSTER* collection or similar container management program site. The cap should not be replaced, but may be taken separately. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

For refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Scarlet Trio

ADVANCED INSECTICIDE

ACTIVE CONSTITUENTS: 116 g/L CLOTHIANIDIN
83 g/L CLOFENTEZINE
13 g/L ABAMECTIN

GROUP 4A 10A 6 INSECTICIDE

For the control of billbug, African black beetle, Argentinian scarab, Argentine stem weevil, couch fly, couch mite, ground pearl, mole cricket in turf as specified in the Directions for Use table.

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.

Indigo Specialty Products Pty Ltd
ABN 15 631 459 660
3/49 Donaldson Road, Rocklea QLD 4106



Indigo Specialty Products ABN 15 631 459 660
3/49 Donaldson Road, Rocklea QLD 4106
Ph: 0402 735 887